Alfonso Aguirre-Muñoz, Donald A. Croll, C. Josh Donlan, R. William Henry III, Miguel Angel Hermosillo, Gregg R. Howald, Bradford S. Keitt, Luciana Luna-Mendoza, Marlenne Rodríguez-Malagón, Luz María Salas-Flores, Araceli Samaniego-Herrera, Jose Angel Sanchez-Pacheco, Jacob Sheppard, Bernie R. Tershy, Jorge Toro-Benito, Shaye Wolf and Bill Wood

High-impact Conservation: Invasive Mammal Eradications from the Islands of Western México

Islands harbor a disproportionate amount of the earth's biodiversity, but a significant portion has been lost due in large part to the impacts of invasive mammals. Fortunately, invasive mammals can be routinely removed from islands, providing a powerful tool to prevent extinctions and restore ecosystems. Given that invasive mammals are still present on more than 80% of the world's major islands groups and remain a premier threat to the earth's biodiversity, it is important to disseminate replicable, scaleable models to eradicate invasive mammals from islands. We report on a successful model from western México during the past decade. A collaborative effort between nongovernmental organizations, academic biologists, Mexican government agencies, and local individuals has resulted in major restoration efforts in three island archipelagos. Forty-two populations of invasive mammals have been eradicated from 26 islands. For a cost of USD 21 615 per colony and USD 49 370 per taxon, 201 seabird colonies and 88 endemic terrestrial taxa have been protected, respectively. These conservation successes are a result of an operational model with three main components: i) a tri-national collaboration that integrates research, prioritization, financing, public education, policy work, capacity building, conservation action, monitoring, and evaluation; ii) proactive and dedicated natural resource management agencies; and iii) effective partnerships with academic researchers in México and the United States. What is now needed is a detailed plan to eradicate invasive mammals from the remaining islands in the region that integrates the needed additional financing, capacity, technical advances, and policy issues. Island conservation in western México provides an effective approach that can be readily applied to other archipelagos where conservation efforts have been limited.

INTRODUCTION

Although islands make up a small percentage of the earth's total area, they support a disproportionate share of global biodiversity (1). Unfortunately, islands have also suffered disproportionately in terms of extinction (2–4). For example, before the onset of anthropogenic impacts ca. 3000 years ago the islands of Oceania supported more than 2000 now extinct bird species—more than 20% of the extant avifauna worldwide (5). Invasive mammals are the primary driver of biodiversity loss on islands and a limited subset of feral mammals are responsible for a majority of insular diversity declines, including rats (*Rattus* spp.), cats (*Felis catus*), goats (*Capra hircus*), pigs (*Sus scrofa*), donkeys (*Equus asinus*), and European rabbits (*Oryctolagus cuniculus*) (6–8).

The 250+ islands of western México show patterns similar to those of islands worldwide (Fig. 1). They are well known for their high biodiversity, endemism, important seabird colonies, and relatively low levels of human disturbance (9–13). Historically, they have been protected from most human perturbations by aridity, isolation, and low human population densities on the adjacent mainland (14). Today, with the exception of the Baja California islands off the Pacific coast, which are in the process of being protected, all the islands of México are legally protected from many forms of land conversion (15, 16). Nonetheless, invasive mammals—primarily rats, cats, goats, pigs, donkeys, and rabbits—have been introduced to many of the islands of western México, causing population declines, a number of extinctions and extirpations, and significant damage to native vegetation (17–23).

Fortunately, invasive mammals can be permanently eradicated from islands. Eradication techniques have been developed in New Zealand and elsewhere during the past three decades, providing a powerful conservation tool (24). During the past decade, a collaborative effort between conservation nongovernmental organizations (NGOs), Mexican government agencies, and local individuals has resulted in the eradication of invasive mammals from the Pacific coast of Baja California, Gulf of California, and Revillagigedos island archipelagos in western México. For these islands we summarize: i) biodiversity patterns, ii) eradication successes, iii) the costs and benefits of those conservation efforts, iv) the general approach used to achieve those conservation gains, and v) the future of island conservation in this region. Island conservation in western México provides a model that may be applicable to other archipelagos. Given that invasive mammals are present on more than 80% of the world's major islands groups and continue to threaten a significant percentage of the earth's biodiversity (25, 26), it is important to disseminate potentially replicable, scaleable models to eradicate invasive mammals from islands.

ISLAND BIODIVERSITY PATTERNS

Islands make up approximately $\sim 3\%$ of land area worldwide, yet they harbor a relatively high percentage of biodiversity, including many endangered species. Of the mammal and bird species that are listed by the World Conservation Union (IUCN) worldwide, 83% of the documented historical extinctions have occurred on islands and 44% of the currently threatened taxa are insular (Fig. 2a). In México, biodiversity patterns are similar: islands make up a mere 0.2% of the total land area, but have suffered at least 50% of México's historic extinctions and hold 18% of the country's endangered mammals and birds (Fig. 2b). Virtually all of the extinctions and extirpations on islands in western México can be attributed to the impact of invasive mammals (18, 20, 22, 23). Because of




Figure 1. The islands of western México where invasive mammals have been successfully removed.




Figure 2. (A) Global percent comparisons of islands (black) versus continental (white) for total area, and extinct and endangered species of mammals and birds combined. (B) Percent comparisons within México between islands (black) and continental (white) for total area, extinct (EX), and endangered (EN) mammals, and extinct and endangered birds. Species data for both figures are from IUCN Red List (www.redlist.org). We classed Red List categories EX and EW as extinct, and CR, EN, and VU as endangered. México area data from Instituto Nacional de Geografía y Estadística.

Table 1. Invasive mammal eradications in western México and the biodiversity protected. Invasive mammals have been removed from 26 islands in the region (BR = black rats, C = cats, D = donkeys, DG = dogs, G = goats, H = horses, M = house mice, Rab = European rabbits).

| Island | Area (ha) | Date | Non-native mammal | | | Endemic Species/Subspecies | | | | Seabird |
|-----------------------------|--------------|------|----------------------|----------|---------|----------------------------|----------|-------|--------|----------------------|
| | | | Removals | Failures | Present | Mammals | Reptiles | Birds | Plants | breeding populations |
| Baja Pacific | | | | | | | | | | |
| Todos Santos North | 62 | 1999 | C,Rab,D ¹ | | | 1 | 0 | 0 | 0 | 7 |
| San Jerónimo | 67 | 2000 | С | | | 1 | 0 | 0 | 0 | 8 |
| Asuncion | 68 | 1994 | С | | | 0 | 0 | 0 | 0 | 9 |
| San Roque | 79 | 1995 | C,BR | | | 0 | 0 | 0 | 0 | 12 |
| Coronado North | 79 | 1995 | С | | | 1 | 4 | 1 | 1 | 10 |
| San Benito Middle | 105 | 1998 | Rab | | | 0 | 0 | 1 | 1 | 15 |
| Todos Santos South | 127 | 1998 | C,Rab | | | 1 | 2 | 0 | 0 | 7 |
| San Benito East | 195 | 2001 | Rab | | | 0 | 0 | 2 | 3 | 15 |
| Coronado South ² | 227 | 2003 | C,G,D ¹ | | | 1 | 2 | 1 | 4 | 8 |
| San Martin | 298 | 2000 | С | | | 1 | 3 | 0 | 0 | 8 |
| San Benito West | 545 | 1998 | Rab,G,D1 | | | 0 | 1 | 2 | 5 | 14 |
| Natividad | 1029 | 1999 | C,G,1 S1 | | | 1 | 0 | 0 | 3 | 10 |
| Gaudalupe | 26 469 | 2005 | Rab, D, H, DG | | G,3 C | 0 | 0 | 4 | 34 | 9 |
| Gulf of California | | | | | | | | | | |
| San Jorge West | 7 | 2000 | BR | | | 0 | 0 | 0 | 0 | 9 |
| San Jorge East | 9 | 2000 | BR | | | 0 | 0 | 0 | 0 | 9 |
| San Jorge Middle | 41 | 2000 | BR | | | 0 | 0 | 0 | 0 | 11 |
| Rasa ⁴ | 59 | 1995 | BR, M | | | 0 | 0 | 0 | 0 | 4 |
| Estanque | 105 | 1999 | C | | | 0 | 1 | 0 | 0 | 2 |
| Isabela ⁵ | 274 | 1996 | С | BR | | 0 | 0 | 0 | 0 | 11 |
| Mejia | 328 | 2001 | С | | | 0 | 1 | 0 | 0 | 5 |
| San Francisco | 465 | 2000 | C, G | | | 2 | 0 | 0 | 0 | 1 |
| Coronados | 1003 | 1999 | C | | | 2 | 2 | 0 | 0 | 1 |
| Monserrate | 1882 | 2001 | С | | | 1 | 2 | 0 | 0 | 2 |
| La Partida | 2029 | 2001 | C | | | 6 | 0 | 0 | 0 | 4 |
| Santa Catalina | 3080 | 2004 | C | | | 1 | 8 | 0 | 1 | 2 |
| Revillagigedo | | | | | | | | | | |
| Clarion | 2928 | 2002 | S,P | Rab | | 0 | 2 | 4 | 13 | 8 |
| Total Protected | 41 559 | | | | | 19 | 28 | 11 | 30 | 201 |

¹Small populations were removed live in cooperation with island residents and governmental agencies. ²Project conducted by Gustavo Arnoud of the Centro de Investigaciones Biológicas, Noroeste, with assistance from Island Conservation. ³While goats are ecological extinct, the eradication campaign is still in progress (less than 200 goats remain). ⁴Project conducted by Jesús Ramírez (deceased) of the Instituto de Ecologia, Universidad Nacional Autónoma de México with assistance from Island Conservation. See Rodriquez et al. (36).

those diversity patterns, the islands of western México are considered a national conservation priority (27).

ERADICATION SUCCESSES

Three small islands with important seabird colonies were the genesis for invasive mammal eradications in México. In 1994/1995, the Mexican NGO Grupo de Ecología y Conservación de Islas (GECI) and the US-based NGO Island Conservation (IC), working with Jesús Ramírez, eradicated feral cats from Asunción Island and cats and black rats (*Rattus rattus*) from nearby San Roque Island (28). That same year Jesús Ramírez and colleagues from Universidad Nacional Autónoma de México eradicated black rats and house mice (*Mus musculus*) from Rasa Island, the world's most important nesting site for Heermann's Gulls (*Larus heermanni*) and Elegant Terns (*Sterna elegans*) (13).

From this beginning, a total of 26 invasive mammal populations have been removed from 13 islands off the Pacific coast of Baja California, 14 populations from 12 islands in the Gulf of California, and two populations from one island in the Revillagigedo Archipelago (Fig. 1, Table 1). Invasive predators (rats, cats, dogs) have been removed from 22 islands and invasive herbivores (donkeys, goats, sheep, and rabbits) from 10 islands. These conservation actions have protected 88 endemic populations from the impacts of invasive mammals (Table 1, Fig. 3).

Entire plant communities are now recovering from extensive overgrazing by invasive herbivores (19, 29). For example, on West San Benito Island, the single island endemic succulent *Dudleyi linearis* was near extinction due to overgrazing by European rabbits; it is now recovering following rabbit



Figure 3. Some of the island species protected by the eradication of invasive mammals on islands in western Mexico. Red-billed tropic-bird (*Phaethon aethereus*) on the San Jorge Islands, the endemic San Martin Island alligator lizard (*Elgaria multicarinata ignara*), the endemic San Jeronimo deer mouse (*Peromyscus maniculatus jeroni*), Cassin's auklet (*Ptychoramphus aleuticus australe*) on the San Benitos Islands, the endemic San Benitos *Dudleya linearis*, the endemic Guadalupe Island pine (*Pinus radiata binata*), and the endemic Guadalupe rock daisy (*Perityle incana*).

Table 2. Measuring conservation impact and cost-effectiveness of non-native mammal eradications in western México.

| | USD-2005 |
|--|-------------|
| Total expenditures (1994–2005) ¹ | \$4 361 318 |
| Cost per seabird colony protected | \$22 139 |
| Cost per endemic terrestrial vertebrate protected ² | \$49 560 |
| Cost per hectare protected ³ | \$58 |

¹Includes all expenditures (salaries, equipment, supplies, transportation, office space, policy work, and other costs) for Grupo de Ecología y Conservación de Islas and Island Conservation activities in México, and in the US and Canada in support of Grupo de Ecología y Conservación de Islas' projects. Also includes reported costs for Isabel Island from Rodriquez et al. (36). Does not include Rasa Island. ²Species or subspecies. ³Does not include i) the removal of rabbits, donkeys, and horses from Guadalupe Island, all of which were by-products of other campaigns or ii) Rasa Island (see Table 1).

eradication (Fig. 3) (28). Recent and ongoing goat removal efforts on Guadalupe Island have resulted in significant plant conservation gains—a total of 34 endemic taxa have been protected, including the rediscovery of four species that were thought to be extinct (29, 30). The endemic pine (*Pinus radiata* var. *binata*) and cypress (*Cupressus guadalupensis*) forests, habitats of the endemic Guadalupe junco (*Junco insularis*), are showing a dramatic comeback (Fig. 3). After a century without recruitment, thousands of pine and cypress seedlings are now present. Upon completion of goat removal from Guadalupe Island, the area of island habitat protected by invasive mammal eradications in México will triple (Table 1).

Eradication successes have also resulted in major conservation gains for seabirds in the region; more than 201 breeding colonies have been protected (Table 1). Many of these colonies represent species that breed on three or fewer islands or total less than 10 000 individuals (31–33). Invasive predators (i.e., cats and rats) have extirpated numerous seabird colonies in the region (e.g., Xantus' Murrelet, *Synthliboramphus hypoleucus*) and now that these predators have been removed, seabirds will likely recolonize the islands, as has been documented on islands to the north in California (20, 34).

During the past decade two eradication efforts in the region have failed, and these failures provide important insights for subsequent island conservation endeavors. In both projects, planning and consultation were limited and pre-eradication trials were not undertaken. In hindsight, the planning phase was particularly crucial because of the novel circumstances of the eradication attempts. Clarion Island is a subtropical island where feral pigs, sheep, and rabbits were targeted for removal. The size of the island (2928 ha) was much larger than any previous rabbit eradication campaign. Before Clarion Island, eradication campaigns led by GECI and IC took place on smaller islands where a small number of highly skilled hunter/ trappers used a largely intuitive approach (35). This approach did not transfer well to Clarion, and the resulting underplanning led to insufficient budgeting. Although pigs and sheep were removed, the rabbit eradication campaign was stopped when techniques were not effective enough and funding ran out. Isabel Island is a tropical island where feral cats and black rats were targeted for removal. A team from Universidad Nacional Autónoma de México successfully eradicated cats, but the black rat eradication failed. This failure was likely because of poor timing of the eradication campaign in relation to rat reproductive biology, hasty project implementation, and unanticipated levels of bait uptake by native land crabs (Coenobita compressus), which are known to consume rodenticides (36). Bait uptake by land crabs, which are immune to most rodenticides, greatly complicates rat eradications on tropical islands and has been responsible for at least two other failures (37, Howald et al. unpubl. data). The failures of these two campaigns highlight the need for i) strategic planning, including consultation and in some cases pre-eradication trials (38), and ii) flexibility to revise project needs when appropriate, such as methods, management, monitoring, evaluation, and budgets.

ERADICATION COSTS AND BENEFITS

The eradication of 42 invasive mammal populations from 26 islands in western México at a cost ~USD4.3 million resulted in high conservation returns on investment (Table 2). Invasive mammals were removed at an average cost of US\$58 per hectare for each invasive species removed. Due to the cost effectiveness of this biodiversity protection and because many species co-occur in México and the United States (33), US government and private foundations have recently taken a trans-border approach to seabird conservation off the Pacific Coast of North America, funding eradication projects in México and, more recently, Canada (39, 40).

CONSERVATION APPROACH

In the for-profit sector, organizations that specialize in a few complex operations often have an efficiency advantage over less specialized organizations (41). Such patterns are also likely to apply to the nonprofit sector, including conservation organizations (42, 43), particularly when highly specialized tasks such as eradicating invasive animals are involved. By specializing in eradications and applying a standardized, integrated, and heterarchial approach, GECI, IC, and Island Conservation-Canada (IC-Canada) have been able to increase efficiency and decrease costs with resulting gains for biodiversity. Perhaps more importantly, the steady accretion of shared expertise across three countries has enabled success in increasingly complex projects (37, 44).

The described conservation successes are a result of an operational model with three main components:

- a tri-national collaboration between three conservation NGOs (GECI, IC, IC-Canada) dedicated to eradicating invasive mammals from islands and preventing new introductions. These three groups applied a common approach that integrated i) research and prioritization, ii) public education, iii) policy work, iv) capacity building, v) on-theground conservation action, and vi) monitoring and evaluation;
- proactive and dedicated government agency partners in México; and
- effective partnerships with academic researchers in México and the United States.

GECI worked closely with these partners and its sister organizations (IC and IC-Canada) to leverage North Americanwide economic, intellectual, and political resources to effectively fund and support science-driven conservation actions. Key stakeholders and agencies include i) universities (Universidad Nacional Autónoma de México, Centro de Investigaciones Biológicas del Noroeste, Centro de Investigación Científica y Educación Superior de Ensenada, University of California Santa Cruz); ii) government agencies (Secretaría de Medio Ambiente y Recursos Naturales, Secretaría de Gobernación, Secretaría de Marina, Armada de México, Instituto Nacional de Ecología, Comisión Nacional de Áreas Naturales Protegidas, Comisión Nacional de Biodiversidad, Procuraduría Federal del Medio Ambiente); and iii) local island users and community fishing co-operatives (Pescadores Nacionales de Abulón, Buzos y Pescadores de Baja California, California de San Ignacio, Abuloneros y Langosteros). This consortium initiated their efforts on smaller islands and used these experiences to build capacity and leverage successes for eradication campaigns on larger, more challenging islands.

Without buy-in from local island residents and users, the benefits of eradication campaigns are potentially short-lived because of the likelihood of intentional or unintentional reintroductions. Thus, many of the eradication campaigns in western México included environmental education campaigns with local fishermen and school children (see 45, 46). Whereas animal rights issues have caused challenges to eradications elsewhere (8), they were not a major issue for the eradications reported here. Environmental education campaigns that focused on the value of native biodiversity and the impacts of invasive mammals may have contributed to the lack of animal rights issues (45).

While the environmental education campaigns facilitate "bottom-up" support, working closely with governmental agencies provides "top-down" support from an island conservation policy perspective. Grupo de Ecología y Conservación de Islas is working with national government agencies to harbor more political support for eradication campaigns and protection of the region's islands in general (47, 48). Both are critical to enable eradication campaigns and prevent reintroductions of invasive species in the long-term.

México has an existing legal infrastructure that affords protection for much of its insular natural heritage. Threatened species are protected by Mexican federal law under the Norma Oficial Mexicana NOM-059-ECOL-2001 (49), with specific protection provided by Article 87 of the General Law of the Ecological Balance and Environmental Protection and Article 85 of the General Wildlife Law. The islands of the Gulf of California have enjoyed legal protection since 1978, and have recently gained World Natural Heritage recognition by United Nations Educational, Scientific and Cultural Organization (16). The protected status of islands off the Pacific side of Baja California is rapidly changing. In 1988, Natividad, San Roque, and Asunción Islands were added to the Vizcaino Biosphere Reserve under the National System of Protected Areas. In 2005, Guadalupe Island and its surrounding islets and waters were designated as the Guadalupe Island Biosphere Reserve as a result of an initiative led by the Comisión Nacional de Areas Naturales Protegidas in partnership with GECI. Currently, the Mexican Federal Congress has mandated a Biosphere Reserve designation for the remaining Pacific Islands. With sufficient resources, these new protected areas could provide both the legal framework and infrastructure for developing island management plans and facilitate the necessary enforcement of existing regulations to protect the insular biodiversity of the entire archipelago. Unfortunately, a lack of resources has inhibited the development of management plans for other protected islands.

Finally, a unique partnership between the IC and University of California Santa Cruz has fostered an integrated approach that combines on-the-ground science and conservation action. Island Conservation is physically housed at University of California's Center for Ocean Health, which facilitates collaborations with academic staff and students (graduate and undergraduate) as well as other academic resources. Grupo de Ecología y Conservación de Islas, based in Ensenada, Baja California, is adopting a similar approach, building relationships with key Mexican academic institutions including Centro de Investigacion Cientifica y de Educacion Superior de Ensenada and Universidad Autónoma de Baja California. Whereas the process of targeting islands for eradications encompasses components that are opportunistic in nature, academic collaborations have helped facilitate a more systematic approach. For example, a regional island biodiversity database was developed to help prioritize eradication efforts in western México (28). Academic analyses of biodiversity patterns and conservation investments have highlighted previously

unrecognized conservation opportunities (33). Concurrent with eradication campaigns, targeted research on select islands also documents the biodiversity impacts of invasive mammals on the region's island ecosystems and the conservation gains of eradication campaigns (18-21, 32, 44, 50). Such research not only helps guide decision-making for future conservation actions, but also aids in fundraising.

THE FUTURE OF ISLAND CONSERVATION IN **WESTERN MÉXICO**

The islands of western México are celebrated laboratories of evolution and important centers of biodiversity (10-13, 51-54). Because of the efforts of multiple government, NGO, and academic institutions, the ability to remove invasive mammals from islands has vastly improved (24, 55-58), contributing to the island conservation gains in México outlined above. Invasive mammals can now be removed from large islands swiftly and cost-effectively (e.g., 59). All of the remaining islands in western México are now feasible conservation targets with respect to invasive mammal eradications. Goats are in the final phases of being removed from Guadalupe Island, one of México's endemic hotspots. However, cats remain on Guadalupe, as well as on other islands with high biodiversity value, such as Socorro Island and the three largest islands in the Islas Marias Archipelago (60–62). Recent advances in techniques and technology make cat eradication on these large islands feasible (63, 64). Invasive rodent populations are still present on eight islands in the region, but many are within the size range where eradication would be straightforward and cost-effective with large conservation returns (58). Funding, rather than technical capacity and island size, are now the limiting factors in preventing extinctions on these and the remaining islands in western México (55).

Fortunately, local, national, and international support for the conservation of México's islands is gaining momentum. What is now needed is a detailed, staged plan to eradicate invasive mammals from the remaining ~60 islands that integrates the needed financing, additional capacity, technical advances, and policy issues. With such a plan and the continued vision and determination of Mexican conservationists, México can continue to demonstrate feasible and cost-effective approaches to restore and safeguard the biodiversity of an entire suite of island archipelagos.

References and Notes

- Myers, N., Mittermeier, R.A., Mittermeier, C.G., Fonseca, G.A.B.d. and Kent, J. 2000. Biodiversity hotspots for conservation priorities. *Nature 403*, 853–858.
- Alcover, J.A., Sans, A. and Palmer, M. 1998. The extent of extinctions of mammals on islands. *J. Biogeogr.* 25, 913–918.
- islands. J. Biogeogr. 25, 913–918.

 Groombridge, B. and Jenkins, M.D. 2002. World Atlas of Biodiversity. Earth's Living Resources in the 21st Century. University of California Press, Berkeley, 256 pp. Martin, P.S. and Steadman, D.W. 1999. Prehistoric extinctions on islands and continents. In: Extinctions in Near Time: Causes, Contexts, and Consequences. R.D.E., MacPhee (ed). Kluwer Academic, New York, pp. 17–53.

 Steadman, D.W. 1995. Prehistoric extinctions of Pacific island birds: biodiversity meets zooarchaeology. Science 267, 1123–1131.

 Atkinson, I.A.E. 1989. Introduced animals and extinctions. In: Conservation for the Treative Extra Contexts. Western, D. and Pearl, M.C. (eds.) Oxford University Press.
- Twenty-First Century. Western, D. and Pearl, M.C. (eds). Oxford University Press, Oxford, pp. 54–75.
 Ebenhard, T. 1988. Introduced birds and mammals and their ecological effects. Swedish
- Wildlife Research Viltrevy 13, 1–107.
 Towns, D.R., Atkinson, I.A.E. and Daugherty, C.H. 2006. Have the harmful effects of
- introduced rats on islands been exaggerated? *Biological Invasions 8*, 863–891. Alvarez-Castañeda, S.T and Patton, J.L. 1999. *Mamiferos del noroeste de Mexico*. Centro de Investigaciones Biologicas del Noroeste, S. C., La Paz, México, 556 pp. Bourillón, L., Cantú, A., Eccardi, F., Lira, E., Ramírez, J., Velarde, E. and González,
- A.Z. 1988. Islas del Golfo de Calfornia. Secretaria de Gobernación, México City, México, 292 pp.
 Case, T.J., Cody, M.L. and Ezcurra, E. 2002. A New Island Biogeography of the Sea of
- Cortes. Oxford University Press, New York, 690 pp.
 Grismer, L.L. 2002. Amphibians and Reptiles of Baja California, Including Its Pacific Islands and the Islands in the Sea of Cortes. University of California Press, Berkeley, 413
- Velarde, E. and Anderson, D.W. 1994. Conservation and management of seabird islands in the Gulf of California: setbacks and successes. In: Seabirds on Islands: Threats, Case Studies and Action Plans. Nettleship, D.N., Burger, J. and Gochfeld, M. (eds). Birdlife International, Cambridge, pp. 229–243.

- Tershy, B.R., Breese, D. and Croll, D.A. 1997. Human perturbations and conservation strategies for San Pedro Martir Island, Islas del Golfo de California Reserve, Mexico. Environ. Conservat. 24, 261–270.
 Carabias-Lillo, J., Maza-Elvira, J.d.I., Gutierrez-Carbonell, D., Gomez-Cruz, M., Anaya-Reyna, G., Zavala-Gonzalez, A., Figueroa, A.L. and Bernudez-Almada, B. 2000. Programa de Manejo Area de Protection de Flora y Fauna Islas de Golfo de California, Mariae Considerable Normal Reservador. Mexico. Comision Nacional de Areas Naturales Protegidas, México City, México, 120
- Ezcurra, E., Bourillón, L., Cantú, A., Martínez, M.E. and Robles, A. 2002. Ecological Conservation. In: A New Island Biogeography of the Sea of Cortés. Case, T.J., Cody,
- M.L. and Ezcurra, E. (eds). Oxford University Press, Oxford, pp. 417–444. Alvarez-Castañeda, S.T. and Ortega-Rubio, A. 2003. Current status of rodents on islands in the Gulf of California. *Biol. Conservat.* 109, 157–163. Donlan, C.J., Knowlton, J., Doak, D.F. and Biavaschi, N. 2005. Nested communities,
- invasive species and Holocene extinctions: evaluating the power of a potential conservation tool. *Oecologia 145*, 475-485.
- Donlan, C.J., Tershy, B.R. and Croll, D.A. 2002. Islands and introduced herbivores:
- conservation action as ecosystem experimentation. *J. Appl. Ecol.* 39, 235–246. Keitt, B.S. 2005. Status of Xantus' murrelet and its nesting habitat in Baja California, Mexico. *Marine Ornithology* 33, 105–114.
- Keitt, B.S., Wilcox, C., Tershy, B.R., Croll, D.A. and Donlan, C.J. 2002. The effect of
- feral cats on the population viability of Black-vented Shearwaters (Puffinus opisthomelas) on Natividad Island, Mexico. Anim. Conservat. 5, 217–223.

 McChesney, G.J. and Tershy, B.R. 1998. History and status of introduced mammals and impacts to breeding seabirds on the California Channel and northwestern Baja California Islands. Colonial Waterbirds 21, 335–347. 22.
- Moran, R. 1996. The flora of Guadalupe Island, Mexico. Memoir. Calif. Acad. Sci. 19, 23.
- Veitch, C.R. and Clout, M.R. (eds). 2002. Turning the Tide: the Eradication of Invasive
- Species. World Conservation Union, Gland, Switzerland, 414 pp.
 Atkinson, I.A.E. 1985. The spread of commensal species of *Rattus* to oceanic islands and Arkinson, 17.2, 1793. It is spread to committee species of Island Birds: Case Studies for the Management of Threatened Island Species. International Council for Bird Preservation No. 3. Moors, P.J. (ed). Cambridge, pp. 35–81.

 Long, J.L. 2003. Introduced Manumals of the World. CSIRO Publishing, Collingwood,
- 26.
- Ceballos, G., Rodriguez, P. and Medellin, R.A. 1998. Assessing conservation priorities in megadiverse Mexico: mammalian diversity, endemicity, and endangerment. Ecol. Appl. 8, 8-17
- Donlan, C.J., Tershy, B.R., Keitt, B.S., Wood, B., Sanchez, J.A., Weinstein, A., Croll, D. and A and Alguilar, J.L. 2000. Island conservation action in northwest Mexico. In: *Proceedings of the Fifth California Islands Symposium*. Browne, D.H., Chaney, H. and Mitchell, K. (eds). Santa Barbara Museum of Natural History, Santa Barbara, pp.
- Keitt, B., Junak, S., Mendoza, L.L. and Aquirre, A. 2005. The restoration of Guadalupe Island. *Fremontia* 33, 20–25.
- Aguirre-Muñoz, A., Ezcurra, E., Enkerlin-H, E., Soberón-M., J., Salas-F., L.M., Santos del Prado, K., Peters, E., Luna-M., L., Tershy, B., García-G., C. and Aguirre-B, I. 2006. La construcción social de la conservación y el desarrollo sustentable de isla Guadalup In: *Isla Guadalupe, Restauración y Conservación*. Santos, K. and Peters, E. (eds). SEMARNAT, INE, CICESE, GECI y SEMAR, Mexico City, pp. 239–254. Donlan, C.J., Howald, G.R., Tershy, B.R. and Croll, D.A. 2003b. Evaluating alternative
- rodenticides for island conservation: roof rat eradication from the San Jorge Islands, Mexico. Biol. Conservat. 14, 29-34.
- Mexico. Biol. Conserval. 14, 29–34.
 Keitt, B.S. and Tershy, B.R. 2003. Cat eradication significantly decreases shearwater mortality. Anim. Conservat. 6, 307–308.
 Wolf, S., Keitt, B., Aquirre, A., Tershy, B., Palacios, E. and Croll, D. 2006. Transboundary seabird conservation in an important North American marine ecoregion. Environ. Conservat. 33, 294–305.
 Whitworth D.L. Carley H.P. Young P. I. Koepke J.S. Grees, E. and Engagon, S.
- ecoregion. Environ. Conservat. 33, 294–305.

 Whitworth, D.L., Carter, H.R., Young, R.J., Koepke, J.S., Gress, F. and Fangman, S. 2005. Initial recovery of Xantus's Murrelets following rat eradication on Anacapa Island, California. Marine Ornithology 33, 131–137.

 Wood, B., Tershy, B.R., Hermosillo, M.A., Donlan, C.J., Sanchez, J.A., Keitt, B.S., Croll, D.A., Howald, G.R. and Biavaschi, N. 2002. Removing cats from islands in northwest Mexico. In: Turning the Tide: The Eradication of Invasive Species. Veitch, C.R. and Clout, M.N. (eds). IUCN SSC Invasive Species Specialist Group, Gland, Switzerland, pp. 374–380.

 Rodríguez, C., Torres, R. and Drummond, H. 2006. Eradicating introduced mammals from a forested tropical island. Biol. Conservat. 130, 98–105.
- from a forested tropical island. Biol. Conservat. 130, 98-105.
- Buckelew, S., Howald, G.R., Wegman, A., Shepard, J., Curl, J., McClelland, P., Tershy,
- Be, Swift, K., et al. 2006. Progress in Palmyra Atoll Restoration: Rat Eradication Trial 2005. Technical Report, Island Conservation, 75 pp.
 Cromarty, P.L., Broome, K.G., Cox, A., Empson, R.A., Hutchinson, W.M. and McFadden, I. 2002. Eradication planning for invasive alien animal species on islands— 38.
- McFadden, I. 2002. Eradication planning for invasive aften animal species on islands—the approach developed by the New Zealand Department of Conservation. In: Turning the tide: the Eradication of Invasive Species. Veitch, C.R. and Clout, M.N. (eds). IUCN SSC Invasive Species Specialist Group, IUCN, Gland, pp. 85–91.

 Luckenbach Trustee Council. 2006. S.S. Jacob Luckenbach and Associated Mystery Oil Spills Draft Damage Assessment and Restoration Plan/Environmental Assessment. Prepared by the California Department of Fish and Game, National Oceanic and Atmospheric Administration, United States Fish and Wildlife Service, National Park Service. (http://www.dfg.ca.gov/ospr/organizational/scientific/nrda/Luckenbach% 20DARP%20combined.ndf) 20DARP%20combined.pdf)
 Montrose Settlements Restoration Program. 2005. Final Restoration Plan and
- Programmatic Environmental Impact Statement, and Environmental Impact Report. Report of the Montrose settlements restoration program, National Oceanic and Atmospheric Administration, US Fish and Wildlife Service, National Park Service, California Department of Fish and Game, California Department of Parks and Recreation, and California State Lands Commission. (http://www.montroserestoration.

- 41. Collins, J. 2001. Good to Great: Why Some Companies Make the Leap and Others Don't.

- Collins, J. 2001. Good to Great: Why Some Companies Make the Leap and Others Don't. Harper Collins, New York, 318 pp. Barr, D.A. 1995. The effects of organizational structure on primary care outcomes under managed care. Ann. Intern. Med. 122, 353–359.

 Roemer, G.W. and Donlan, C.J. 2005. Biology, policy and law in endangered species conservation: II. a case history in adaptive management of the island fox on Santa Catalina Island, California. Endangered Species Update 22, 144–156.

 Howald, G.R., Faulkner, K.R., Tershy, B., Keitt, B., Gellerman, H., Creel, E.M., Grinnel, M., Ortega, S., et al. 2005. Eradication of black rat from Anacapa Island: biological and social considerations. In: Proceedings of the Sixth California Islands Symposium. Garcelon, D.K., and Schwemm, C.A. (eds). Institute for Wildlife Studies. Symposium. Garcelon, D.K. and Schwemm, C.A. (eds). Institute for Wildlife Studies,
- Arcata, pp. 299–312.

 Donlan, C.J. and Keitt, B.S. 1999. Using research and education to prevent extinction.
- California Coast and Ocean 15, 20–23.
 Tershy, B.R., Donlan, C.J., Keitt, B., Croll, D., Sanchez, J.A., Wood, B., Hermosillo, M.A. and Howald, G. 2002. Island conservation in northwest Mexico: a conservation
- model integrating research, education and exotic mammal eradication. In: Turning the Tide: the Eradication of Invasive Species. Veitch, C.R. and Clout, M.N. (eds). IUCN SSC Invasive Species Specialist Group. IUCN, Gland, pp. 293–300. Aguirre-Muñoz, A., Samaniego-Herrera, A., Garcia Gutiérrez, C., Enkerlin-Hoeflich, E., Carranza-Sánchez, J., Sánchez-Ibarra, C., Luna-Mendoza, L.M., Keitt, B., et al. 2005. Propuesta para el establecimiento del Area Natural Protegida Reserva de la Biosfera Islanda Deputition. Estudio Técnico Instituto Grupo de Escologia y Conservación de Islanda Deputition.
- Islas del Pacifico. Estudio Técnico Justificativo. Grupo de Ecología y Conservación de Islas, A.C. Ensenada, Baja California, México, 126 pp.
 Aguirre-Muñoz, A., B.-C. J, Carranza, J., Enkerlin-Hoeflich, E., García-Gutiérrez, C., Luna-Mendoza, L.M., Keitt, B., Sanchéz-Pacheco, J.A. and Tershy, B.R. 2003. Propuesta para el establecimiento del Area Natural Protegida Reserva de la Biósfera de la Isla Guadalupe. Estudio Técnico Justificativo. Grupo de Ecología y Conservación de
- Islas, A.C. Ensenada, Baja California, México, 78 pp.
 Secretaría de medio ambiente y recursos naturales. 2002. Norma Oficial Mexicana NOM-059-2001. In: Protección ambiental-Especies nativas de México de Flora y Fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo. Publicado en el D.O.F. de fecha 06 de marzo de 2002, Máxico. DE pp. 14. México, D.F, pp. 1–4.
 Donlan, C.J., Croll, D.A. and Tershy, B.R. 2003a. Islands, exotic herbivores and

- Donlan, C.J., Croll, D.A. and Tershy, B.R. 2003a. Islands, exotic herbivores and invasive plants: their roles in coastal California conservation. *Restor. Ecol. 11*, 524–530. Avise, J.C., Smith, M.H., Selander, R.K., Lawlor, T.E. and Ramsey, P.R. 1974. Biochemical polymorphism and systematics in the genus Peromyscus. V. insular and mainland species of the subgenus Haplomylomys. *Sytematic Zoology 23*, 226–238. Felger, R.S. and Lowe, C.H. 1976. The island and coastal vegetation and flora of the northern part of the Gulf of California. *Natural History Museum of Los Angeles County Contributions in Science 285*, 1–58.

 Santos del Prado, K. and Peters, E. (eds). 2006. *Isla Guadalupe, restauración y*
- Santos del Prado, K. and Peters, E. (eds). 2006. Ista Guadatupe, restauración y conservación. Instituto Nacional de Ecologia, México City.
 Soule, M. and Yang, S.Y. 1972. Genetic variation in side-blotched lizards on island in the Gulf of California. Evolution 27, 593–600.
 Campbell, K. and Donlan, C.J. 2005. Feral goat eradications on islands. Conservat. Biol.
- 19, 1362–1374.
- 19, 1502–1574.

 Donlan, C.J., Tershy, B.R., Campbell, K. and Cruz, F. 2003c. Research for requiems: the need for more collaborative action in invasive species management and conservation.

 Conservat. Biol. 17, 1850–1851.

 Lavoie, C., Donlan, C.J., Campbell, K., Cruz, F. and Carrion-G., V. 2007. Geographic
- tools for eradication programs of insular non-native mammals. Biological Invasions 9,
- Towns, D.R. and Broome, K.G. 2003. From small Maria to massive Campbell: forty
- years of rat eradications from New Zealand islands. New Zeal. J. Zool. 30, 377–398. Cruz, F., Donlan, C.J., Campbell, K. and Carrion, V. 2005. Conservation action in the Galápagos: Feral pig (Sus scrofa) eradication from Santiago Island. Biol. Conservat. 121,
- Keitt, B., Henry, R.W., Aquirre, A., Garcia, C., Mendoza, L.L., Hermosillo, M.A, Tershy, B. and Croll, D. 2006. Impacts of introduced cats (*Felis catus*) on the Guadalupe Island ecosystem. In: *Taller sobre la restauración y conservación de Isla Guadalupe*: memorias. d. Prado, G.K.S. and Peters, E. (eds). Instituto Nacional de Ecología, México, D.F., pp. 15-24.
- Martínez-Gómez, J.E. and Curry, R.L. 1996. The conservation status of the Socorro Mockingbird Minodes graysoni in 1993–94. Bird Conservation International 6, 271–283. Rodriguez-Estrella, R., Moreno, A.R. and Tam, K.G. 2000. Spring diet of the endemic
- ring-tailed cat (Bassariscus astutus insulicola) population on an island in the Gulf of California, Mexico. J. Arid Environ, 44, 241–246.
- Algar, D. and Burrows, N.D. 2004. Feral cat control research: Western Shield review-February 2003. Conservation Science Western Australia 5, 131–163. Phillips, R.B., Cooke, B.D., Campbell, K., Carrion, V., Marquez, C. and Snell, H.L.
- 2005. Eradicating feral cats to protect Galapagos land iguanas: methods and strategies. Pacific Conservation Biology 11, 57–66.

 The authors' names are listed in alphabetical order. The work presented here is the result of a decade of collaborative work. We thank our conservation partners mentioned in the manuscript, as well the Vizcaino Biosphere Reserve, Islas del Golfo de California Reserve, Bahia Loreto Marine Park, Grupo Ecologista Antares, Centro de Estudio del Designto y Oceano, Pro-Esteros, Pro-Natura, and Nijagaria, We thank the many island Desierto y Oceano, Pro-Esteros, Pro-Natura, and Niparaja. We thank the many island users who opened up their islands and their homes to us and helped us in the field. Teunding was provided by International Coalition for Bird Preservation, USFWS International, Conservation International, Packard Foundation, National Fish & Wildlife Foundation, USFWS Region 1&2, Weeden Foundation, Conservation Food & Health Foundation, Switzer Foundation, Blank Family Foundation, US National Park Service, WWF–Mexico, Fundo Mexicano para la Conservación de la Naturaleza, Oracle Corporation, Walton Foundation, Farallon Island Foundation, Seacology, the Sandler Family Foundation, and an important anonymous supporter. This work would Sandler Family Foundation, and an important anonymous supporter. This work would not have been possible without the assistance of J. and V. Davis, D. Seymore, E. and R. Tershy, V. McDermit, A. Acevedo and B. Bedolfe. B. Bell, A. Saunders, and D. Veitch provided valuable advice.
- First submitted 4 January 2007. Accepted for publication 2 July 2007.

Alfonso Aguirre-Muñoz is the director of Grupo de Ecología y Conservación de Islas. His address: Grupo de Ecología y Conservación de Islas AC, Avenida López Mateos No. 1590-3, Ensenada, Baja California, México, CP 22880.

E-mail: alfonso.aguirre@conservaciondeislas.org

Donald Croll is a co-founder of Island Conservation (IC) and a professor of Ecology and Evolutionary Biology at the University of California Santa Cruz (UCSC). His address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA, or University of California Santa Cruz, Department of Ecology and Evolutionary Biology, Long Marine Lab, 100 Shaffer Road, Santa Cruz, CA 95060, USA. E-mail: croll@ucsc.edu

Josh Donlan is the founder and director of Advanced Conservation Strategies and has worked with IC during the past decade. His address: Advanced Conservation Strategies, PO Box 8254, Santa Cruz, CA 95061, USA, and Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY 14853, USA.

E-mail: cjd34@cornell.edu

R. William Henry III is a PhD candidate at UCSC and a biologist with IC. His address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA, and University of California Santa Cruz, Department of Ecology and Evolutionary Biology, Long Marine Lab, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail: rhenry@ucsc.edu

Miguel Angel Hermosillo is a conservation technician with Grupo de Ecología y Conservación de Islas. His address: Grupo de Ecología y Conservación de Islas AC, Avenida López Mateos No. 1590-3, Ensenada, Baja California, México, CP 22880. E-mail: miguel.hermosillo@conservaciondeislas.org

Gregg R. Howald is the founder and director of Island Conservation-Canada. His address: Island Conservation Canada, 680–220 Cambie Street, Vancouver, British Columbia, V6B 2M9, Canada.

 $\hbox{E-mail: ghowald@islandconservation.org}\\$

Bradford C. Keitt is program director with IC. His address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail: brad.keitt@islandconservation.org

Luciana Luna-Mendoza is a biologist with Grupo de Ecología y Conservación de Islas. Her address: Grupo de Ecología y Conservación de Islas AC, Avenida López Mateos No. 1590-3, Ensenada, Baja California, México, CP 22880.

E-mail: Luciana.Mendoza@conservaciondeislas.org

Marlenne Rodríguez-Malagón is a biologist with Grupo de

Ecología y Conservación de Islas. Her address: Grupo de Ecología y Conservación de Islas AC, Avenida López Mateos No. 1590-3, Ensenada, Baja California, México, CP 22880. E-mail: Marlenne.Rodriguez@conservaciondeislas.org

Luz María Salas-Flores is the Oceanography Research Coordinator for the Mexican Navy. Her address: Secretaría de Marina-Armada de México, Segunda Zona Naval, Ensenada, Mexico, BC.

E-mail: luz@govt.mx

Araceli Samaniego-Herrera is a biologist with Grupo de Ecología y Conservación de Islas. Her address: Grupo de Ecología y Conservación de Islas AC, Avenida López Mateos No. 1590-3, Ensenada, Baja California, México, CP 22880.

E-mail: Araceli.Samaniego@conservaciondeislas.org

Jose Angel Sanchez-Pacheco is the founder and former director of Grupo de Ecología y Conservación de Islas. His address: Instituto SummerTree AC, Calle Diorita #271. Colonia Pedregal Playitas, Ensenada, Baja California, México, CP 22840. E-mail: jasanpa2000@hotmail.com

Jacob Shepard is a biologist with IC. His address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail: jacob.shephard@islandconservation.org

Bernie Tershy is a co-founder and executive director of IC and an adjunct professor in Ecology and Evolutionary Biology at UCSC. His address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail: tershy@islandconservation.org

Jorge Toro-Benito is the sub-director of Insular Territories Management, Secretariat of the Interior for the Mexican federal government. His address: Subdirección Administración del Territorio Insular, Unidad de Gobierno, Secretaría de Gobernación, Rio Amazonas No. 62 - 38 Piso, Col. Cuauhtémoc, México, DE 06500.

E-mail: benito@govt.mx

Shaye Wolf is a PhD candidate at UCSC and biologist with IC. Her address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA, and University of California Santa Cruz, Department of Ecology and Evolutionary Biology, Long Marine Lab, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail: wolf@biology.ucsc.edu

Bill Wood is a conservation specialist with IC. His address: Island Conservation, Center for Ocean Health, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail: wood@islandconservation.org